



Urinary Pain Relief[®] Maximum Strength

AZO

Quick Reference Guide

Product Name	AZO Urinary Pain Relief [®] Maximum Strength, 97.5 mg
Indications and Usage	AZO Urinary Pain Relief [®] Maximum Strength contains Phenazopyridine Hydrochloride, an orally administered urinary tract analgesic. Phenazopyridine HCl is indicated for the symptomatic relief of pain, burning, urgency, frequency arising from irritation of the lower urinary tract mucosa caused by infection, trauma, surgery, or medical procedures. Phenazopyridine HCl provides only symptom relief and should be used as directed on the package.
Dosage and Administration	Take 2 tablets, 3 times daily with or after meals as needed for up to two days.
Storage and Handling	Store at room temperature (59°– 86°F) Store in a cool, dry place and protect from light
Clinical Profile	Phenazopyridine HCl is prescribed for the symptomatic relief of the pain, burning, urgency and frequency typically experienced with urinary tract infections. ¹ Phenazopyridine HCl has been widely used in the United States since 1914 for this purpose. Phenazopyridine HCl is excreted in the urine (65% of the oral dosage is chemically unchanged) where it exerts a topical analgesic effect on the mucosa of the urinary tract. ^{1,2} The benefits are provided within 1 hour, the time in which Phenazopyridine HCl reaches the bladder as indicated by a change in urine color. ³
How Supplied	Supplied in dark red/brown tablet (size and color may vary) containing 97.5mg of Phenazopyridine HCl. Tablets are supplied in blisters as a 12 count or 24 count.

¹Hardman, J.G., L.E. Limbird, P.B. Molinoff, R.W. Ruddon, A.G. Goodman (eds.). Goodman and Gilman's The Pharmacological Basis of Therapeutics. 9th ed. New York, NY: McGraw-Hill, 1996. p. 1070.

²Warner Chilcott - Pyridium Rx (http://www.wcrx.com/pdfs/pi/pyridium_pi_cpc.pdf).

³Hui, Jane Y C, Marie-Andrée Harvey, and Shawna L Johnston. "Confirmation of ureteric patency during cystoscopy using phenazopyridine HCl: a low-cost approach." Journal of obstetrics and gynaecology Canada 31.9 (2009) 845-849.

FOR PROFESSIONAL USE ONLY